arXiv:1603.00681v2 [math.NT] 21 Mar 2016

The Bi-periodic Fibonacci Octonions

Nazmiye Yilmaz*
Department of Mathematics, Science Faculty,
Selcuk University, Campus, Konya, Turkey

Yasin Yazlik
Department of Mathematics, Faculty of Science and Letters,
Nevsehir Haci Bektas Veli University, Nevsehir, Turkey

Necati Taskara
Department of Mathematics, Science Faculty,
Selcuk University, Campus, Konya, Turkey

March 22, 2016

Abstract

In this paper, by using bi-periodic Fibonacci numbers, we introduce the bi-periodic Fibonacci octonions. After that, we derive the generating function of these octonions as well as investigated some properties over them. Also, as another main result of this paper, we give the summations for bi-periodic Fibonacci octonions.

Keywords: Bi-periodic Fibonacci sequence, Fibonacci sequence, generating functions, octonions.
Ams Classification: 11B39, 11R52, 15A66, 20G20.

1 Introduction

There has been an increasing interest on quaternions and octonions that play an important role in various areas such as computer sciences, physics, differential geometry, quantum physics, signal, color image processing, geostatics and analysis [1, 8].

Let $\mathcal{O}$ be the octonion algebra over the real number field $\mathbb{R}$. It is known, by the Cayley-Dickson process that any $p \in \mathcal{O}$ can be written as

$$p = p' + p''e\,,$$

*e mail: *nzyilmaz@selcuk.edu.tr, yyazlik@nevsehir.edu.tr, ntaskara@selcuk.edu.tr*

where $p', p'' \in H = \left\{ a_0 + a_1 i + a_2 j + a_3 k \,\middle|\, i^2 = j^2 = k^2 = -1, ijk = -1,\ a_0, a_1, a_2, a_3 \in \mathbb{R} \right\}$ the real quaternion division algebra. The octonions in Clifford algebra C are a normed division algebra with eight dimensions over the real numbers larger than the quaternions. The field $\mathcal{O} \cong C^4$ of octonions

$$p = \sum_{s=0}^{7} p_s e_s, \quad p_s \in \mathbb{R},$$

is an eight-dimensional non-commutative and non-associative $\mathbb{R}$-field generated by eight base elements

$$e_0 = 1,\ e_1 = i,\ e_2 = j,\ e_3 = k,\ e_4 = e,\ e_5 = ie,\ e_6 = je,\ e_7 = ke\,.$$

The addition and multiplication of any two octonions, $p = p' + p''e$, $q = q' + q''e$, are defined by

$$p + q = p' + q' + (p'' + q'')e \ \text{ and } \ pq = p'q' - \overline{q''}p'' + (q''p' + p''\overline{q'})e\,,$$

where $\overline{q'}, \overline{q''}$ denote the conjugates of the quaternions q', q'', respectively. The conjugate and norm of p are denoted by

$$\overline{p} = Re(p) - Im(p) \text{ and } p\overline{p} = \sum_{s=0}^{7} p_s^2\,, \tag{1.1}$$

respectively [1, 8, 38, 42].

On the other hand, the literature includes many papers dealing with the special number sequences such as Fibonacci, Lucas, Pell ([4, 9, 21, 25, 26, 30, 39, 40, 41]). One of these directions goes through to the *bi-perodic Fibonacci* (or, equivalently, *generalized Fibonacci*) and the *bi-periodic Lucas* (or, equivalently, *generalized Lucas*). In fact bi-perodic Fibonacci sequences have been firstly defined by Edson and Yayenie in 2009 and then some important properties for these sequences have been investigated in the references [9, 39]. Also, bi-perodic Lucas sequences have been defined by Bilgici in 2014 and then some important properties for these sequences have been investigated in the reference [4]. For any two nonzero real numbers a and b, the bi-periodic Fibonacci $\{q_n\}_{n=0}^{\infty}$ and bi-periodic Lucas sequences $\{l_n\}_{n=0}^{\infty}$ are defined recursively by

$$q_0 = 0,\ q_1 = 1,\ q_n = \begin{cases} aq_{n-1} + q_{n-2}, & \text{if } n \text{ is even} \\ bq_{n-1} + q_{n-2}, & \text{if } n \text{ is odd} \end{cases} \qquad n \geq 2 \tag{1.2}$$

and

$$l_0 = 2,\ l_1 = a,\ l_n = \begin{cases} bl_{n-1} + l_{n-2}, & \text{if } n \text{ is even} \\ al_{n-1} + l_{n-2}, & \text{if } n \text{ is odd} \end{cases} \qquad n \geq 2\,. \tag{1.3}$$

In addition, the authors in the references [4],[9] and [39] expressed so many properties on the bi-periodic Fibonacci and bi-periodic Lucas sequences. In fact some of the main outcomes (that depicted in these references) of these sequences can be summarized as in the following:

- The Binet formulas are given by

$$q_n = \frac{a^{1-\xi(n)}}{(ab)^{\lfloor \frac{n}{2} \rfloor}} \left(\frac{\alpha^n - \beta^n}{\alpha - \beta} \right) \tag{1.4}$$

and

$$l_n = \frac{a^{\xi(n)}}{(ab)^{\left\lfloor \frac{n+1}{2} \right\rfloor}} \left(\alpha^n + \beta^n\right), \tag{1.5}$$

where the condition $\xi(n) = n - 2\left\lfloor \frac{n}{2} \right\rfloor$ can be read as

$$\xi(n) = \begin{cases} 0, & n \text{ is even} \\ 1, & n \text{ is odd} \end{cases} \tag{1.6}$$

and α, β are roots of characteristic equation of $\lambda^2 - ab\lambda - ab = 0$.

- The generating functions for the bi-periodic Fibonacci and bi-periodic Lucas sequences with odd subscripted are

$$f(x) = \sum_{m=1}^{\infty} q_{2m-1} x^{2m-1} \frac{x - x^3}{1 - (ab+2)x^2 + x^4}, \tag{1.7}$$

and

$$L_1(x) = \sum_{m=0}^{\infty} l_{2m+1} x^{2m+1} \frac{a + ax^3}{1 - (ab+2)x^2 + x^4}. \tag{1.8}$$

- The bi-perodic Fibonacci and bi-peridoic Lucas sequences satify the equations

$$(ab+4)q_n = l_{n-1} + l_{n+1}, \tag{1.9}$$

$$l_n = q_{n-1} + q_{n+1} \tag{1.10}$$

and

$$q_{-n} = (-1)^{n-1} q_n, \tag{1.11}$$

$$l_{-n} = (-1)^n l_n. \tag{1.12}$$

After all these above material on bi-periodicity, let us give our attention to the other classification parameters on these above special sequences, namely quaternions and octonions. We should note that, in the literature (see [2, 3, 5]-[7], [10]-[20],[22]-[24],[27]-[29], [31]-[37] and the references cited in them), it can be found some works on the quaternions and octonions for Jabosthal, Pell as well as the Fibonacci and Lucas numbers. For example, in [24], it is investigated some properties of the Fibonacci and Lucas octonions.

It is natural to wonder whether there exits a connection between the parameters bi-periodicity and quaternions. Actually this was done in [35, 36]. The authors defined the bi-periodic Fibonacci and Lucas quaternions as

$$Q_n = \sum_{s=0}^{3} q_{n+s} e_s \text{ and } P_n = \sum_{s=0}^{3} l_{n+s} e_s,$$

where $n \geq 0$.

In the light of all these above material (depicted as separate paragraphs), the main goal of this paper is to define *bi-periodic Fibonacci octonions* with a different viewpoint. To do that, by using bi-periodic Fibonacci numbers, we obtain the generating function of these octonions. We also actually investigated some properties of bi-periodic Fibonacci octonions.

2 The bi-periodic Fibonacci octonions

As we indicated in the previous section, in this part, we introduce the bi-periodic Fibonacci octonions as a new generalization of Fibonacci octonions by considering the bi-periodic Fibonacci numbers.

Definition 2.1 *The bi-periodic Fibonacci octonions $O_n(a,b)$ are defined by*

$$O_n(a,b) = \sum_{s=0}^{7} q_{n+s} e_s, \tag{2.1}$$

where q_n is the bi-periodic Fibonacci number for $n \in \mathbb{N} = \{0,1,2,3,\ldots\}$.

From the Equation (1.11), that is $q_{-n} = (-1)^{n-1} q_n$, the bi-periodic Fibonacci octonions with negative subscripts are defined by

$$O_{-n}(a,b) = \sum_{s=0}^{7} (-1)^{n-s-1} q_{n-s} e_s, \tag{2.2}$$

where $n \in \mathbb{N} = \{0,1,2,3,\ldots\}$.

After all, we present the following first proposition of this paper.

Proposition 2.1 *For $n \in \mathbb{N}$, we give the results related to bi-periodic Fibonacci octonions:*

i) $O_n(a,b) + \overline{O_n(a,b)} = 2q_n e_0$,

ii) $O_n^2(a,b) + O_n(a,b)\overline{O_n(a,b)} = 2q_n e_0 O_n(a,b)$,

iii) $O_n(a,b)\overline{O_n(a,b)} = \sum_{s=0}^{7} q_{n+s}^2$

iv) $O_{-n}(a,b) + \overline{O_{-n}(a,b)} = 2(-1)^{n-1} q_n e_0$

v) $O_n(a,b) + O_{-n}(a,b) = l_n O_0(a,b)$, *where n is even.*

Proof.

i) From Definition 2.1, we obtain

$$\begin{aligned} O_n(a,b) + \overline{O_n(a,b)} &= \sum_{s=0}^{7} q_{n+s} e_s + q_n e_0 - \sum_{s=1}^{7} q_{n+s} e_s \\ &= 2q_n e_0 \,. \end{aligned}$$

ii) By taking into account *i)*, we have

$$\begin{aligned} O_n^2(a,b) &= O_n(a,b)(2q_n e_0 - \overline{O_n(a,b)}) \\ &= 2q_n e_0 O_n(a,b) - O_n(a,b)\overline{O_n(a,b)} \,. \end{aligned}$$

iii) The proof is easily seen by using the Equation (1.1).

iv) The proof is similar to *i)*.

v) From the Equations (2.1) and (2.2), we write for n even

$$\begin{aligned} O_n(a,b)+O_{-n}(a,b) &= \sum_{s=0}^{7} q_{n+s}e_s + \sum_{s=0}^{7}(-1)^{n-s-1}q_{n-s}e_s \\ &= \sum_{s=1}^{7}(q_{n+s}+(-1)^{s-1}q_{n-s})e_s\,. \end{aligned}$$

By taking into account the Equations (1.2) and (1.10), we obtain

$$\begin{aligned} O_n(a,b)+O_{-n}(a,b) &= l_n\sum_{s=1}^{7} q_s e_s \\ &= l_n O_0(a,b)\,. \end{aligned}$$

■

Theorem 2.1 *For $n \geq 0$, the Binet formula for the bi-periodic Fibonacci octonions is*

$$O_n(a,b)=\begin{cases} \dfrac{1}{(ab)^{\lfloor \frac{n}{2} \rfloor}}\left(\dfrac{\alpha^*\alpha^n-\beta^*\beta^n}{\alpha-\beta}\right), & \text{if } n \text{ is even} \\ \dfrac{1}{(ab)^{\lfloor \frac{n}{2} \rfloor}}\left(\dfrac{\alpha^{**}\alpha^n-\beta^{**}\beta^n}{\alpha-\beta}\right), & \text{if } n \text{ is odd} \end{cases}$$

where α,β are roots of characteristic equation of $\lambda^2-ab\lambda-ab=0$ and
$\alpha^*=\sum_{t=0}^{7}\dfrac{a^{\xi(t+1)}}{(ab)^{\lfloor \frac{t}{2} \rfloor}}\alpha^t e_t$ *and* $\beta^*=\sum_{t=0}^{7}\dfrac{a^{\xi(t+1)}}{(ab)^{\lfloor \frac{t}{2} \rfloor}}\beta^t e_t$,
$\alpha^{**}=\sum_{t=0}^{7}\dfrac{a^{\xi(t)}}{(ab)^{\lfloor \frac{t+1}{2} \rfloor}}\alpha^t e_t$ *and* $\beta^{**}=\sum_{t=0}^{7}\dfrac{a^{\xi(t)}}{(ab)^{\lfloor \frac{t+1}{2} \rfloor}}\beta^t e_t$.

Proof. It can easily seen by considering the Definition 2.1 and the Equation (1.4). ■

Theorem 2.2 *For $n,r \in \mathbb{Z}^+$ and $n \geq r$, the following equality is hold:*

$$O_{2n-2r}(a,b)O_{2n+2r}(a,b)-O_{2n}^2(a,b)=\frac{\alpha^*\beta^*((ab)^{2r}-\alpha^{4r})+\beta^*\alpha^*((ab)^{2r}-\beta^{4r})}{(ab)^{2r}(\alpha-\beta)^2}.$$

Proof. Let us take $U=O_{2n-2r}(a,b)O_{2n+2r}(a,b)-O_{2n}^2(a,b)$. Then, by using Teorem 2.1, we write

$$\begin{aligned} U &= \frac{1}{(ab)^{\lfloor \frac{2n-2r}{2} \rfloor}}\left(\frac{\alpha^*\alpha^{2n-2r}-\beta^*\beta^{2n-2r}}{\alpha-\beta}\right)\frac{1}{(ab)^{\lfloor \frac{2n+2r}{2} \rfloor}}\left(\frac{\alpha^*\alpha^{2n+2r}-\beta^*\beta^{2n+2r}}{\alpha-\beta}\right) \\ &\quad -\left(\frac{1}{(ab)^{\lfloor \frac{2n}{2} \rfloor}}\left(\frac{\alpha^*\alpha^{2n}-\beta^*\beta^{2n}}{\alpha-\beta}\right)\right)^2 \\ &= \frac{1}{(ab)^{2n}(\alpha-\beta)^2}[\alpha^*\beta^*(-ab)^{2n}(1-\alpha^{4r}(-ab)^{-2r})+\beta^*\alpha^*(-ab)^{2n}(1-\beta^{4r}(-ab)^{-2r})] \\ &= \frac{\alpha^*\beta^*((ab)^{2r}-\alpha^{4r})+\beta^*\alpha^*((ab)^{2r}-\beta^{4r})}{(ab)^{2r}(\alpha-\beta)^2}\,. \end{aligned}$$

■

If we take $r = 1$ in Theorem 2.2, it is obtained in the following corollary.

Corollary 2.1 *For $n \in \mathbb{Z}^+$, we have the Cassini-like identity*

$$O_{2n-2}(a,b)O_{2n+2}(a,b) - O_{2n}^2(a,b) = \frac{\alpha^*\beta^*((ab)^2 - \alpha^4) + \beta^*\alpha^*((ab)^2 - \beta^4)}{(ab)^2(\alpha-\beta)^2}.$$

Theorem 2.3 *For $n, r \in \mathbb{Z}^+$, $n \geq r$ and r is even, we have*

$$O_{n-r}(a,b)O_{n+r}(a,b) - O_n^2(a,b) = \begin{cases} \frac{\alpha^*\beta^*((ab)^r - \alpha^{2r}) + \beta^*\alpha^*((ab)^r - \beta^{2r})}{(ab)^r(\alpha-\beta)^2}, & \text{if } n \text{ is even} \\ -\frac{\alpha^{**}\beta^{**}((ab)^r - \alpha^{2r}) + \beta^{**}\alpha^{**}((ab)^r - \beta^{2r})}{(ab)^{r-1}(\alpha-\beta)^2}, & \text{if } n \text{ is odd} \end{cases}.$$

Proof. The proof is done similarly to the proof of Theorem 2.2. ■

Now, we present the generating function of the bi-periodic Fibonacci octonions in the following theorem.

Theorem 2.4 *For the bi-periodic Fibonacci octonions, we have the generating function*

$$\sum_{n=0}^{\infty} O_n(a,b)x^n = \frac{O_0(a,b) + x(O_1(a,b) - bO_0(a,b)) + (a-b)R(x)}{1 - bx - x^2},$$

where

$R(x) = (xe_0 + e_1 + \frac{1}{x}e_2 + \frac{1}{x^2}e_3 + \frac{1}{x^3}e_4 + \frac{1}{x^4}e_5 + + \frac{1}{x^5}e_6 + \frac{1}{x^6}e_7)f(x) - (xe_1 + e_2 + (\frac{1}{x} + (ab+1)x)e_3 + (\frac{1}{x^2} + ab + 1)e_4 + (\frac{1}{x^3} + (ab+1)\frac{1}{x} + (a^2b^2 + 3ab + 1)x)e_5 + (\frac{1}{x^4} + (ab+1)\frac{1}{x^2} + (a^2b^2 + 3ab + 1))e_6 + (\frac{1}{x^5} + (ab+1)\frac{1}{x^3} + (a^2b^2 + 3ab + 1)\frac{1}{x} + (a^3b^3 + 5a^2b^2 + 6ab + 1)x)e_7)$

and $f(x) = \dfrac{x - x^3}{1 - (ab+2)x^2 + x^4}$.

Proof. Assume that $G(x,a,b)$ is the generating function for the bi-periodic Fibonacci octonions. Then we have

$$G(x,a,b) = O_0(a,b) + O_1(a,b)x + O_2(a,b)x^2 + \cdots + O_n(a,b)x^n + \cdots. \tag{2.3}$$

If it is multiplying equation (2.3) with bx and x^2, respectively, then we have

$$bxG(x,a,b) = bO_0(a,b)x + bO_1(a,b)x^2 + bO_2(a,b)x^3 + \cdots + bO_n(a,b)x^{n+1} + \cdots. \tag{2.4}$$

$$x^2G(x,a,b) = O_0(a,b)x^2 + O_1(a,b)x^3 + O_2(a,b)x^4 + \cdots + O_n(a,b)x^{n+2} + \cdots. \tag{2.5}$$

By considering the Equations (1.2), (2.3), (2.4), (2.5) and Definition 2.1, it is obtained the equation

$$\begin{aligned} (1 - bx - x^2)G(x) &= O_0(a,b) + x(O_1(a,b) - bO_0(a,b)) \\ &\quad + \sum_{n=2}^{\infty}(O_n(a,b) - bO_{n-1}(a,b) - O_{n-2}(a,b))x^n \\ &= O_0(a,b) + x(O_1(a,b) - bO_0(a,b)) \\ &\quad + \sum_{n=2}^{\infty}\sum_{s=0}^{7}(q_{n+s} - bq_{n+s-1} - q_{n+s-2})e_s x^n. \end{aligned}$$

By considering again the Equations (1.2) and (1.7), we get

$$\begin{aligned}(1-bx-x^2)G(x) &= O_0(a,b)+x(O_1(a,b)-bO_0(a,b))+(a-b)\sum_{n=1}^{\infty}q_{2n-1}x^{2n}e_0\\ &+(a-b)\sum_{n=2}^{\infty}q_{2n-1}x^{2n-1}e_1+(a-b)\sum_{n=2}^{\infty}q_{2n-1}x^{2n-2}e_2\\ &+(a-b)\sum_{n=3}^{\infty}q_{2n-1}x^{2n-3}e_3+(a-b)\sum_{n=3}^{\infty}q_{2n-1}x^{2n-4}e_4\\ &+(a-b)\sum_{n=4}^{\infty}q_{2n-1}x^{2n-5}e_5+(a-b)\sum_{n=4}^{\infty}q_{2n-1}x^{2n-6}e_6\\ &+(a-b)\sum_{n=5}^{\infty}q_{2n-1}x^{2n-7}e_7\,.\end{aligned}$$

Consequently, we have

$$\begin{aligned}(1-bx-x^2)G(x) &= O_0(a,b)+x(O_1(a,b)-bO_0(a,b))+(a-b)xf(x)e_0\\ &+(a-b)(f(x)-q_1x)e_1+(a-b)x^{-1}(f(x)-q_1x)e_2\\ &+(a-b)x^{-2}(f(x)-q_1x-q_3x^3)e_3\\ &+(a-b)x^{-3}(f(x)-q_1x-q_3x^3)e_4\\ &+(a-b)x^{-4}(f(x)-q_1x-q_3x^3-q_5x^5)e_5\\ &+(a-b)x^{-5}(f(x)-q_1x-q_3x^3-q_5x^5)e_6\\ &+(a-b)x^{-6}(f(x)-q_1x-q_3x^3-q_5x^5-q_7x^7)e_7\\ &= O_0(a,b)+x(O_1(a,b)-bO_0(a,b))+(a-b)R(x)\end{aligned}$$

as required. ■

For bi-periodic Fibonacci octonions, we give the summations according to specified rules.

Theorem 2.5 *For $n\in\mathbb{Z}^+$, there exist*

i) $\sum_{r=0}^{n-1}O_r(a,b)=\dfrac{O_{n+1}(a,b)+O_n(a,b)-O_{n-1}(a,b)-O_{n-2}(a,b)}{ab}$
$+\dfrac{\alpha^*\beta-\beta^*\alpha-\alpha^{**}ab+\beta^{**}ab}{ab(\alpha-\beta)}$,

ii) $\sum_{r=0}^{n-1}O_{2r}(a,b)=\dfrac{O_{2n}(a,b)-O_{2n-2}(a,b)}{ab}+\dfrac{\alpha^*\beta-\beta^*\alpha}{ab(\alpha-\beta)}$,

iii) $\sum_{r=0}^{n-1}O_{2r+1}(a,b)=\dfrac{O_{2n+1}(a,b)-O_{2n-1}(a,b)}{ab}-\dfrac{\alpha^{**}-\beta^{**}}{\alpha-\beta}$.

Proof. We will, now, just prove *i*), since the proofs of *ii*) and *iii*) can be done quite similarly with it. The main point of the proof will be touched just the result Theorem 2.1(in other words the Binet formula of these octonions). We must note that the proof should be investigated for both cases of n.

$\underline{n \textbf{ is odd }}$**:** In this case, we get

$$\begin{aligned}
\sum_{r=0}^{n-1} O_r(a,b) &= \sum_{r=0}^{\frac{n-1}{2}} O_{2r}(a,b) + \sum_{r=0}^{\frac{n-3}{2}} O_{2r+1}(a,b) \\
&= \sum_{r=0}^{\frac{n-1}{2}} \frac{1}{(ab)^r}\left(\frac{\alpha^*\alpha^{2r} - \beta^*\beta^{2r}}{\alpha-\beta}\right) + \sum_{r=0}^{\frac{n-3}{2}} \frac{1}{(ab)^r}\left(\frac{\alpha^{**}\alpha^{2r+1} - \beta^{**}\beta^{2r+1}}{\alpha-\beta}\right) \\
&= \frac{\alpha^*}{\alpha-\beta}\sum_{r=0}^{\frac{n-1}{2}}\left(\frac{\alpha^2}{ab}\right)^r - \frac{\beta^*}{\alpha-\beta}\sum_{r=0}^{\frac{n-1}{2}}\left(\frac{\beta^2}{ab}\right)^r + \frac{\alpha^{**}}{\alpha-\beta}\sum_{r=0}^{\frac{n-3}{2}}\alpha\left(\frac{\alpha^2}{ab}\right)^r \\
&\quad -\frac{\beta^{**}}{\alpha-\beta}\sum_{r=0}^{\frac{n-3}{2}}\beta\left(\frac{\beta^2}{ab}\right)^r
\end{aligned}$$

$$\begin{aligned}
\sum_{r=0}^{n-1} O_r(a,b) &= \frac{\alpha^*}{\alpha-\beta}\frac{\left(\frac{\alpha^2}{ab}\right)^{\frac{n-1}{2}+1}-1}{\frac{\alpha^2}{ab}-1} - \frac{\beta^*}{\alpha-\beta}\frac{\left(\frac{\beta^2}{ab}\right)^{\frac{n-1}{2}+1}-1}{\frac{\beta^2}{ab}-1} \\
&\quad + \frac{\alpha^{**}}{\alpha-\beta}\frac{\left(\frac{\alpha^2}{ab}\right)^{\frac{n-3}{2}+1}\alpha-\alpha}{\frac{\alpha^2}{ab}-1} - \frac{\beta^{**}}{\alpha-\beta}\frac{\left(\frac{\beta^2}{ab}\right)^{\frac{n-3}{2}+1}\beta-\beta}{\frac{\beta^2}{ab}-1}.
\end{aligned}$$

In here, simplifying the last equality, we imply the equality in i) as required.

$\underline{n \textbf{ is even }}$**:** From Theorem 2.1, we know

$$\begin{aligned}
\sum_{r=0}^{n-1} O_r(a,b) &= \sum_{r=0}^{\frac{n-2}{2}} O_{2r}(a,b) + \sum_{r=0}^{\frac{n-2}{2}} O_{2r+1}(a,b) \\
&= \sum_{r=0}^{\frac{n-2}{2}} \frac{1}{(ab)^r}\left(\frac{\alpha^*\alpha^{2r} - \beta^*\beta^{2r}}{\alpha-\beta}\right) + \sum_{r=0}^{\frac{n-2}{2}} \frac{1}{(ab)^r}\left(\frac{\alpha^{**}\alpha^{2r+1} - \beta^{**}\beta^{2r+1}}{\alpha-\beta}\right) \\
&= \frac{\alpha^*}{\alpha-\beta}\frac{\left(\frac{\alpha^2}{ab}\right)^{\frac{n-2}{2}+1}-1}{\frac{\alpha^2}{ab}-1} - \frac{\beta^*}{\alpha-\beta}\frac{\left(\frac{\beta^2}{ab}\right)^{\frac{n-2}{2}+1}-1}{\frac{\beta^2}{ab}-1} \\
&\quad + \frac{\alpha^{**}}{\alpha-\beta}\frac{\left(\frac{\alpha^2}{ab}\right)^{\frac{n-2}{2}+1}\alpha-\alpha}{\frac{\alpha^2}{ab}-1} - \frac{\beta^{**}}{\alpha-\beta}\frac{\left(\frac{\beta^2}{ab}\right)^{\frac{n-2}{2}+1}\beta-\beta}{\frac{\beta^2}{ab}-1}.
\end{aligned}$$

As a result, by arranging the last equality, we have

$$\sum_{r=0}^{n-1} O_r(a,b) = \frac{O_{n+1}(a,b) + O_n(a,b) - O_{n-1}(a,b) - O_{n-2}(a,b)}{ab} + \frac{\alpha^*\beta - \beta^*\alpha - \alpha^{**}ab + \beta^{**}ab}{ab(\alpha-\beta)}.$$

■

Corollary 2.2 *In the all results of Section 2, to reveal the importance of this subject, we can express certain and immediate relationships as follows:*

- *If we replace $a = b = 1$ in q_n, we obtain the same result in [24] for Fibonacci octonions.*
- *If we replace $a = b = 2$ in q_n, we obtain the same result in [34] for Pell octonions.*
- *If we replace $a = b = k$ in q_n, we get the k-Fibonacci octonions.*

Conclusion 2.1 *In this paper, we define the octonions for bi-periodic Fibonacci sequences and present some properties of these octonions. By the results in Sections 2 of this paper, we have a great opportunity to compare and obtain some new properties over these octonions. This is the main aim of this paper. Thus, we extend some recent result in the literature.*

In the future studies on the octonions for number sequences, we except that the following topics will bring a new insight.

- *It would be interesting to study the different summations and properties for bi-periodic Fibonacci octonions,*
- *Also, it would be interesting to investigate the octonions for bi-periodic generalized Fibonacci sequences.*

Acknowledgement 2.1 *This research is supported by TUBITAK and Selcuk University Scientific Research Project Coordinatorship (BAP).*

References

[1] Adler SL.,Quaternionic quantum mechanics and quantum fields, New York: Oxford University Press, 1994.

[2] Akkus I., Kecilioglu O., Split Fibonacci and Lucas Octonions, *Advances in Applied Clifford Algebras*, 25, 2015, 517-525.

[3] Akyigit M., Kosal H.H., Tosun M., Fibonacci generalized quaternions, *Advances in Applied Clifford Algebras*, 24, 2014, 631-641.

[4] Bilgici G., Two generalizations of Lucas sequence, *Applied Mathematics and Computation*, 245, 2014, 526-538.

[5] Catarino P., A note on h(x)-Fibonacci quaternion polynomials, *Chaos Solitons and Fractals*, 77, 2015, 1-5.

[6] Catarino P., The Modified Pell and the Modified k-Pell Quaternions and Octonions, *Advances in Applied Clifford Algebras*, Doi 10.1007/s00006-015-0611-4, 2015.

[7] Cimen CB., Ipek A., On Pell quaternions and Pell-Lucas quaternions, *Advances in Applied Clifford Algebras*, 26, 2016, 39-51.

[8] Dixon G.M., Division Algebras: Octonions, Quaternions, Complex Numbers and the Algebraic Design of Physics, Kluvwer Academic Publishers, ISBN 0-7923-2890-6, 1994.

[9] Edson M., Yayenie O., A new generalization of Fibonacci sequences and extended Binet's Formula, *Integers*, 9(A48), 2009, 639-654.

[10] Flaut C., Savin D., Quaternion Algebras and Generalized Fibonacci-Lucas Quaternions, *Advances in Applied Clifford Algebras*, 25, 2015, 853-862.

[11] Flaut C., Shpakivskyi V., On generalized Fibonacci quaternions and Fibonacci-Narayana quaternions, *Advances in Applied Clifford Algebras*, 23, 2013, 673-688.

[12] Flaut C., Shpakivskyi V., Some remarks about Fibonacci elements in an arbitrary algebra, *Bulletin de la Socit des Sciences et des Lettres de d*, LXV(3), 2015, 63-73.

[13] Halici S., On Dual Fibonacci octonions, *Advances in Applied Clifford Algebras*, 25, 2015, 905-914.

[14] Halici S., On Fibonacci quaternions, *Advances in Applied Clifford Algebras*, 22, 2012, 321-327.

[15] Horadam AF., Complex Fibonacci Numbers and Fibonacci Quaternions, *Amer. Math. Monthly*, 70, 1963, 289-291.

[16] Horadam AF., Quaternion recurrence relations, *Ulam Q.*, 2, 1993, 22-33.

[17] Iakin AL., Extended Binet forms for generalized quaternions of higher order, *Fibonacci Quarterly*, 19, 1981, 410-413.

[18] Iakin AL., Generalized quaternions of higher order,*Fibonacci Quarterly*, 15, 1977, 343-346.

[19] Iakin AL., Generalized quaternions with quaternion components, *Fibonacci Quarterly*, 15, 1977, 350-352.

[20] Ipek A., Ari K., On h(x)-Fibonacci octonion polynomials, *Alabama Journal of Mathematics*, 39, 2015, 1-6.

[21] Irmak N., Alp M., Some identities for generalized Fibonacci and Lucas sequences, *Hacet. J. Math.*, 42, 2013, 331-338.

[22] Iyer MR., A note on Fibonacci quaternions, *Fibonacci Quarterly*, 3, 1969, 225-229.

[23] Iyer MR., Some results on Fibonacci quaternions, *Fibonacci Quarterly*, 7, 1969, 201-210.

[24] Kecilioglu O., Akkus I., The Fibonacci Octonions, *Advances in Applied Clifford Algebras*, 25, 2015, 151-158.

[25] Koshy T., Fibonacci and Lucas Numbers with Applications, John Wiley and Sons Inc, NY, 2001.

[26] Panario D, Sahin M,Wang Q, A family of Fibonacci-like conditional sequences, *Integers*, 13(A78), 2013.

[27] Polatli E., A generalization of Fibonacci and Lucas Quaternions, *Advances in Applied Clifford Algebras*, Doi 10.1007/s00006-015-0626-x, 2015.

[28] Polatli E., Kesim S., On quaternions with generalized Fibonacci and Lucas number components, *Adv. Differ. Equ.*, 2015, 1-8.

[29] Ramirez J., Some combinatorial properties of the k-Fibonacci and the k-Lucas quaternions, *An. St. Univ. Ovidius Constanta Ser. Mat.*, 23(2), 2015, 201-212.

[30] Sahin M., The Gelin-Ces‘aro identity in some conditional sequences, *Hacet. J. Math.*, 40, 2011, 855-861.

[31] Savin D., Some properties of Fibonacci numbers, Fibonacci octonions, and generalized Fibonacci-Lucas octonions, *Advances in Difference Equations*, Doi 10.1186/s13662-015-0627-z, 2015, 1-10.

[32] Swamy MNS., On generalized Fibonacci quaternions, *Fibonacci Quarterly*, 5, 1973, 547-550.

[33] Szynal-Liana A., Włoch I., A Note on Jacobsthal Quaternions, *Advances in Applied Clifford Algebras*, 26, 2016, 441-447.

[34] Szynal-Liana A., Włoch I., The Pell Quaternions and the Pell Octonions, *Advances in Applied Clifford Algebras*, 26, 2016, 435-440.

[35] Tan E., Yilmaz S., Sahin M., A note on bi-periodic Fibonacci and Lucas quaternions, *Chaos, Solitons and Fractals*, 85, 2016, 138-142.

[36] Tan E., Yilmaz S., Sahin M., On a new generalization of Fibonacci quaternions, *Chaos, Solitons and Fractals*, 82, 2016, 1-4.

[37] Tasci D., Yalcin F., Fibonacci-p Quaternions, *Advances in Applied Clifford Algebras*, 25, 2015, 245-254.

[38] Tian Y., Matrix representations of octonions and their applications, *Advances in Applied Clifford Algebras*, 10(1), 2000, 61-90.

[39] Yayenie O., A note on generalized Fibonacci sequence, *Applied Mathematics and Computation*, 217(12), 2011, 5603-5611.

[40] Yazlik Y., Taskara N., A note on generalized k-Horadam sequence, *Computers & Mathematics with Applications*, 63, 2012, 36-41.

[41] Yilmaz N., Yazlik Y., Taskara N., On the k-Generalized Fibonacci Numbers, *Selcuk Journal of Applied Mathematics*, 13(1), 2012, 83-88.

[42] Ward J.P., Quaternions and Cayley Numbers, Kluwer, Dordrecht, 1997.